#### **Global Ice Story**

By Joe D'Aleo, CCM, AMS Fellow Co-Founder The Weather Channel

January 27, 2021

The Arctic includes the Arctic Ocean, Greenland, Iceland, and part of Siberia and northern Alaska. Because of the absence of any land mass in the Arctic Ocean, most of area lacks glaciers, which require a land mass. Thus, most of the Arctic contains only floating sea ice. Greenland, Iceland, northern Alaska, and northern Siberia contain the only glaciers in the general Arctic region.

Arctic warming and the melting of the arctic ice are not at all unprecedented (they happen predictably on multidecadal scales with a period of around 60 years) and are in fact entirely natural. Warming results in part from the reduction of arctic ice extent because of flows of warm water into the arctic from the Pacific through the Bering Straits and from the Atlantic from the far North Atlantic Current. The warmer water thins the ice from beneath, slows the refreezing and limits to varying degrees the depth and extent of the ice.



The arctic ice unlike the Glaciers on land in Greenland, the Antarctic and in the world's mountain ranges is floating on water and the melting has no effect on sea level (much as ice in cold beverages doesn't cause the liquid to overflow the glass when it melts).

The Monthly Weather Review in 1922 alerted to the start of this warming period "a radical change in climatic conditions, and hitherto unheard-of high temperatures"

ftp://ftp.library.noaa.gov/docs.lib/htdocs/rescue/mwr/064/mwr-064-02-c1.pdf

Polyakov et al (2002) created a temperature record using stations north of 62 degrees N. The late 1930s-early 1940s were clearly the warmest of the last century. In addition, the numbers of available observations in the late 1930s-early 1940s (slightly more than 50) is comparable to recent decades.



Arctic Basin wide temperatures (Polyakov 2003)

### MULTIDECADAL OCEAN CYCLES AND THE ARCTIC

The Japan Agency for Marine-Earth Science and Technology in Yokosuka, Kanagawa Prefecture observed in a story in Yahoo Asia News in 2005 an ice shrinkage in the western Arctic Ocean from 1997 to 1998 that they attributed to "... by the flow to the area of warm water from the Pacific Ocean, not by atmospheric impact as previously thought". This was related to the super El Nino of 1997/98. JAMSTEC's Koji Shimada, the group's sub-leader, said the shrinkage was particularly severe in the Pacific side of the Arctic Ocean. The ocean's ratio of area covered with ice during the summer stood at about 60-80 percent from the 1980s to mid-1990s, but it went down to 15-30 percent after 1998, he said. Trenberth (1999) also has acknowledged this warming effect of El Nino on the arctic.

The cycle with respect to arctic temperatures and ice relate to multidecadal cycles in ocean temperatures in both the Pacific (Pacific Decadal Oscillation or PDO) and Atlantic (Atlantic Multidecadal Oscillation or AMO).

#### THE PACIFIC DECADAL OSCILLATION (PDO)

The Pacific warm mode favors more El Ninos and warmer water in the far northern Pacific including the Bering Straits. The PDO flipped into its warm mode in 1978 and the arctic temperatures began to warm and ice began to melt repeating what happened in the 1910s to early 1940s time period.





Notice how the temperatures in Alaska go through step changes tied to the PDO (Keen, 2011).



In the winter 0f 2019/20, at Fairbanks average daytime high was -2.1F, while the daytime low -20F, average daily mean was -11F! The coldest was -43F while the warmest was 31F December 9. 33 days were at or below -30F, 5 were at or below - 40F. The winter averaged 4.7F below normal, the 3<sup>rd</sup> coldest since before the Great Pacific Climate Shift (a shift of the so called Pacific Decadal Oscillation to positive) in the late 1970s when warmer Pacific waters favored warmth in Alaska and western North America.





# THE ATLANTIC'S ROLE

The Atlantic also cycles on a 60-70 year period. The Atlantic Multidecadal Oscillation or AMO returned to the positive warm mode in 1995.



Frances et al. (GRL 2007) showed how the warming in the arctic and the melting ice was related to warm water (+3C) in the Barents Sea moving slowly into the Siberian arctic and melting the ice. She also noted the positive feedback of changed "albedo" due to open water then further enhances the warming.

The International Arctic Research Center at the University of Alaska, Fairbanks showed how arctic temperatures have cycled with intrusions of Atlantic water - cold and warm beneath the ice.



Of the two oceans, for the larger arctic basin, the Atlantic may be more important.

Pryzbylak (2000) says:

"There exists an agreement in estimating temperature tendencies prior to 1950. Practically all (old and new) of the papers which cover this time period concentrate on the analysis of the significant warming which occurred in the Arctic from 1920 to about 1940....In the Arctic, the highest temperatures since the beginning of instrumental observation occurred clearly in the 1930s. Moreover, it has been shown that even in the 1950s the temperature was higher than in the last 10 years."

*"For arctic temperature, the most important factor is a change in the atmospheric circulation over the North Atlantic" The influence of the atmospheric circulation changes over the Pacific (both in the northern end and in the tropical parts) is significantly lower"* 

As is the case for US temperatures, the combination of the PDO and AMO Indexes (PDO+AMO) again has considerable explanatory power for Arctic average temperature.



Arctic basin wide temperatures from Polyakov (2003) versus PDO+AMO (STD). Dark blue is annual and purple 5 year running means.

"In approximately the last 140 years, there have been two periods of significant temperature increases in the Arctic. The first began in around 1918–1920 and lasted until 1938 and has been called the '1930s warming' (Bengtsson et al. 2004). Other works have referred to this period as the 'Early Twentieth Century Warming' (ETCW, Brönnimann 2009) or the 'Early Twentieth Century Arctic Warming' (ETCAW, Wegmann et al. 2017, 2018)

Karlen (2005) reported on historical temperatures in Svalbard (Lufthavn, at 78 deg N latitude), claiming that the area represents a large portion of the Arctic. It is reported that the "mean annual temperature increased rapidly from the 1910s to the late 1930s." Later, temperatures dropped, "and a minimum was reached around 1970." Once again, "Svalbard thereafter became warmer, but the mean temperature in the late 1990s was still slightly cooler than it was in the late 1930s."

Drinkwater (2006) concluded that "in the 1920s and 1930s, there was a dramatic warming of the air and ocean temperatures in the northern North Atlantic and the high Arctic, with the largest changes occurring north of 60°N," which "led to reduced ice cover in the Arctic and subarctic regions and higher sea temperatures." This was "the most significant regime shift experienced in the North Atlantic in the 20th century.".

Hanna, et al (2006) estimated Sea Surface Temperatures (SSTs) near Iceland over a 119-year period based on measurements made at ten coastal stations located between latitudes 63°'N and 67°'N. They concluded that there had been "generally cold conditions during the late nineteenth and early twentieth centuries; strong warming in the 1920s, with peak SSTs typically being attained around 1940; and cooling thereafter until the 1970s, followed once again by warming - but not generally back up to the level of the 1930s/1940s warm period."

# THE EFFECT ON ICE COVER

Both the Atlantic and Pacific play roles in arctic ice extent. The sea ice extent diminished following the Great Pacific Climate Shift (flip of the PDO to positive) in the late 1970s. It stayed relatively stable until early 2000s when a more precipitous decline began, related to a spike in North Atlantic warmth and a positive AO.



Dmitrenko and Polyokov (2003) observed that warm Atlantic water in the early 2000s from the warm AMO that developed in the middle 1990s had made its way under the ice to off of the arctic coast of Siberia where it thinned the ice by 30% much as it did when it happened in the last warm AMO period from the 1880s to 1930s.

The University of Colorado's National Snow and Ice Data Center (NSIDC) summarized the role of the ocean cycles very well in October 2007 in this way:

"One prominent researcher, Igor Polyakov at the University of Fairbanks, Alaska, points out that pulses of unusually warm water have been entering the Arctic Ocean from the Atlantic, which several years later are seen in the ocean north of Siberia. These pulses of water are helping to heat the upper Arctic Ocean, contributing to summer ice melt and helping to reduce winter ice growth.

Another scientist, Koji Shimada of the Japan Agency for Marine–Earth Science and Technology, reports evidence of changes in ocean circulation in the Pacific side of the Arctic Ocean. Through a complex interaction with declining sea ice, warm water entering the Arctic Ocean through Bering Strait in summer is being shunted from the Alaskan coast into the Arctic Ocean, where it fosters further ice loss.

Many questions still remain to be answered, but these changes in ocean circulation may be important keys for understanding the observed loss of Arctic sea ice."

### THE SUN AS THE ULTIMATE DRIVER FOR CLIMATE IN ARCTIC

As to the driver for the multidecadal ocean cycles it may well be the solar cycles. Soon (GRL 2005) showed how the Arctic temperatures (Polyakov) correlated extremely well with the total solar irradiance (Hoyt-Schattem) (r-squared of 0.79). This compared to an r-squared correlation of just 0.22 with the CO2.



#### GREENLAND

Greenland is often considered a sign of climate change. The data here too suggests it is cyclical and relates to ocean cycles on century scales and longer. Note the current interglacial period ice core data tells us the last 10,000 years shows 1000-year cycles. They have slowly declined as we move towards the end of the interglacial period (usually just over 10,000 years in length. The last warm period called the Medieval Warm Period saw ice retreat in Greenland enough to entice the Vikings to settle Greenland and grow crops including grapes for wine. They abandoned it as we cooled towards the Little Ice Age.



Data from Nuuk in western Greenland shows the greatest warming from 1880 to the 1930s with a secondary warming since the 1990s Atlantic warming.



### ANTARCTICA

The figure below shows temperature changes during the past 30 years, compared to temperatures recorded between 1950 and 1980. Most of Antarctica is cooling, with warming occurring over just a small portion of the continent that juts out into the Southern Ocean. That region is volcanic with both surface and sea bottom vents. It is also subject to periodic warming from winds that blow downslope and locally warm by compression.



#### Temperature (K), 1984-2014 minus 1950-1980

# THE ANTARCTIC ICE SHEET IS GROWING, NOT MELTING.

Previous studies showed cooling across all of the much larger East Antarctic Ice Sheet and warming limited to the Antarctic Peninsula of west Antarctica. In 2009, Steig et al. published a controversial paper, "Warming of the Antarctic Ice-Sheet Surface Since the 1957 International Geophysical Year," in Nature contending that warming instead was occurring.



Antarctic temperatures show that Antarctica is cooling, not warming

Measured satellite and surface temperatures confirm the lack of warming over most of Antarctica. The UAH and RSS satellite records are the most comprehensive.



UAH Antarctic satellite temperatures show no warming for 37 years.



RSS Antarctic satellite temperatures show no warming for 37 year



The main conclusion to be drawn from these data is that at least 95% of glacial ice in Antarctica is increasing, not melting.

### Cooling of the Southern Ocean around Antarctica

The Southern Ocean around Antarctica has been getting markedly colder since 2006. Sea ice has increased substantially, especially since 2012.



Temperature anomalies of the Southern Ocean showing sharp cooling since 2006.

# WEST ANTARCTIC ICE SHEET

The West Antarctic Ice Sheet occupies a deep basin west of the main East Antarctic Ice Sheet. It comprises only about 8% of glacial ice in Antarctica. The Antarctic Peninsula has been cooling sharply since 2006. Ocean temperatures have been plummeting since about 2007, sea ice has reached all-time highs, and surface temperatures at 13 stations on or near the Antarctic Peninsula have been cooling since 2000. A plot of temperature anomalies at 13 Antarctic stations on or near the Antarctic Peninsula show that the Antarctic Peninsula was warming up until 2000 but has been cooling dramatically since then.



Temperature anomalies at 13 Antarctic stations on or near the Antarctic Peninsula, showing that the Antarctic Peninsula was warming up until 2000 but has been cooling dramatically since then.

The Larsen Ice Shelf Station has been cooling at an astonishing rate of  $1.8^{\circ}$  C per decade ( $18^{\circ}$ C per century) since 1995 (Fig. 8). Nearby Butler Island records even faster cooling at 1.9 C/decade. Sea ice around Antarctica is increasing because ocean temperature from the surface to 100 m dropped below the freezing point in 2008 and has stayed there since.

Annual average temperature at the Larsen Ice Shelf shows sharp cooling (1.8 ° C/decade). (Easterbrook, 2016) (From GISTemp.)

SUMMARY

Multidecadal Oscillations in the Pacific and the Atlantic are acknowledged to be the result of natural processes. The warm mode of the Pacific results in warm water off Alaska that can enter the arctic through the Bering Strait and produce arctic ice melt. The warm mode of the AMO also results in warming in the North Atlantic waters, which are carried by the North Atlantic current into the arctic reducing ice depth and extent. When you combine the two cycles, you can explain the temperature and ice cover variances of the past 110 years for the Arctic. Greenland data suggests the recent warming falls far short of earlier warming periods during the current interglacial and short of the warming early in the 20<sup>th</sup> century. The Antarctic has cooled and ice has increased in recent years although volcanism near the Antarctic peninsula leads to local water warmth and sea ice melting. Prior to the recent meting, the ice cover reached a long-term record high.

Also we should note that the prescribed melting reported in the Science Journal can't be claimed a long time record as global ocean data prior to the satellite (1980) and Argo Buoy era (post 2004) is spotty at best.

Even if the claims about water released were true, computations show global sea level would rise just 4 inches/century (agreeing with global data) and not the up to 24 feet promised decades ago. See the sea level story <u>here</u>.

#### References:

Arctic Climate Assessment (ACIA), 2004. Impacts of a warming Arctic. Cambridge University Press, Cambridge,UK

Changnon, S., Winstanley, D.:2004: Insights to Key Questions about Climate Change, Illinois State Water Survey http://www.sws.uiuc.edu/pubdoc/IEM/ISWSIEM2004-01.pdf

Christy, J.R., R.W. Spencer and W.D. Braswell, 2000: MSU tropospheric temperatures: Dataset construction and radiosonde comparisons. J. Atmos. Oceanic Tech., 17, 1153-1170.

Climate Change Impacts on the United States: The Potential Consequences of Climate Variability and Change National Assessment Synthesis Team USGRCP, June 2000

Delworth, T.L. ,and M.E. Mann, 2000: Observed and simulated multidecadal variability in the Northern Hemisphere. Climate Dyn., 16, 661–676.

Drinkwater, K.F. 2006. The regime shift of the 1920s and 1930s in the North Atlantic. Progress in Oceanography **68**: 134-151.

Gray, S.T., et al., 2004: A tree-ring based reconstruction of the Atlantic Multidecadal Oscillation since 1567 A.D.Geophys. Res. Lett., 31, L12205, doi:10.1029/2004GL019932

Hanna, E., Jonsson, T., Olafsson, J. and Valdimarsson, H. 2006. Icelandic coastal sea surface temperature records constructed: Putting the pulse on air-sea-climate interactions in the Northern North Atlantic. Part I: Comparison with HadISST1 openocean surface temperatures and preliminary analysis of long-term patterns and anomalies of SSTs around Iceland. Journal of Climate **19**: 5652-5666.

Hass, C., Eicken, H., 2001: Interannual Variability of Summer Sea Ice thickness in the Siberian and central Arctic under Different Atmospheric Circulation Regiomes, JGR, 106, 4449-4462

IPCC Fourth Assessment 2007

Johannessewn, O.M., Shalina, E.V., Miles, M. W., (1999): Satellite Evidence for an Arctic Sea Ice Cover in Transformation, Science, 286, 1937-1939

*Karlen, W. 2005. Recent global warming: An artifact of a too-short temperature record?* Ambio **34**: 263-264.

Kerr, R. A., A North Atlantic climate pacemaker for the centuries, Science, 288 (5473), 984-1986, 2000.

Latif, M. and T.P. Barnett, 1994: Causes of decadal climate variability over the North Pacific and North America. Science 266, 634-637.

Polyakov, I., Walsh, D., Dmitrenko, I., Colony, R.L. and Timokhov, L.A. 2003a. Arctic Ocean variability derived from historical observations. Geophysical Research Letters 30: 10.1029/2002GL016441.

Polyakov, I., Alekseev, G.V., Timokhov, L.A., Bhatt, U.S., Colony, R.L., Simmons, H.L., Walsh, D., Walsh, J.E. and Zakharov, V.F., 2004. Variability of the Intermediate Atlantic Water of the Arctic Ocean over the Last 100 Years. Journal of Climate 17: 4485-4497.

Proshutinsky, A.Y., Johnson, M.A., 1997: Two Circulation Regimes of the Wind Driven Arctic, JGR, 102, 12493-12514

Przybylak, R., 2000, Temporal And Spatial Variation Of Surface Air Temperature Over The Period Of Instrumental Observations In The Arctic, Intl Journal of Climatology, **20**: 587–614

Rigor, I.G., Wallace, J.M. and Colony, R.L., 2002. Response of Sea Ice to the Arctic Oscillation. Journal of Climate 15: 2648-2663.

Rothrock, D.A., Yu, Y., Maykut, G.A., 1999: Thinning of the Arctic Sea-Ice Cover, GRL, 26, no23 3469-3472

Soon, W.H., "Variable Solar Irradiance as a Plausible Agent for Multidecadal Variations the Arctic-wide Surface Air Temperature Record of the Past 130 Years," Geophysical Research Letters, Vol. 32, doi:10.1029/2005GL023429.

Thomas, R., Akins, T., Csatho, B., Fahenstock, M., Goglneni, P., Kim, C., Sonntag, J., (2000): Mass Balance of the Greenland Ice Sheet at High Elevations, Science, 289, 427

Trenberth, K.E., and J.W. Hurrell, 1999: Decadal atmosphere-ocean variations in the Pacific. Clim. Dyn., 9, 303-319.

Venegas, S.A., Mysak, L.A., 2000: Is There a Dominant Time scale of Natural Climate Variability in the Arctic, Journal of Climate, October 2000,13, 3412-3424

Wadhams , P., Davis, N.R., 2000: Further Evidence of Ice thinning in the Arctic Ocean, GRL, 27, 3973-3975

*Winsor, P.,(2001) Arctic Sea ice Thickness Remained Constant During the 1990s: GRL 28, no6 1039-1041*